

**EVALUATION
OF CHEMICAL
COMPOSITION
AND IN VITRO
ANTICHOLINESTERIC
ACTIVITY OF ESSENTIAL
OILS FROM HYMENAEA
CANGACEIRA AND
EUGENIA GRACILLIMA**

Natália Sônia de Sá Gomes Feitosa Rosa

Faculty of Integration of Sertão-FIS, Serra
Talhada-PE, Brazil

<http://lattes.cnpq.br/0320635223403287>

Bruno José da Silva Bezerra

Federal University of Pernambuco,
Department of Biochemistry, Recife-PE,
Brazil

<http://lattes.cnpq.br/5524038913919195>

*Layra Mirelly Feitosa Ferreira Diniz
Ventura*

Faculty of Integration of Sertão-FIS, Serra
Talhada-PE, Brazil

<http://lattes.cnpq.br/3496411924340327>

Janiely Feitosa de Carvalho

Faculty of Integration of Sertão-FIS, Serra
Talhada-PE, Brazil

<http://lattes.cnpq.br/7607148797957841>

Gabriela Pereira Calumbi

Faculty of Integration of Sertão-FIS, Serra
Talhada-PE, Brazil

<http://lattes.cnpq.br/4784625946115118>

All content in this magazine is licensed under a Creative Commons Attribution License. Attribution-Non-Commercial-Non-Derivatives 4.0 International (CC BY-NC-ND 4.0).



Estefani Pontes Simão

Sertão School of Medicine – FMS,
Arcoverde-PE, Brazil
<http://lattes.cnpq.br/5576357171256259>

Bruno José do Nascimento

University of Pernambuco, Department of
Animal Morphology and Physiology, Recife-
PE, Brazil
<http://lattes.cnpq.br/8213260513385508>

Álvaro Aguiar Coelho Teixeira

University of Pernambuco, Department of
Animal Morphology and Physiology, Recife-
PE, Brazil
<http://lattes.cnpq.br/1539131079574469>

Valéria Wanderley Teixeira

University of Pernambuco, Department of
Animal Morphology and Physiology, Recife-
PE, Brazil
<http://lattes.cnpq.br/4292195468804301>

Maria Tereza dos Santos Correia

Federal University of Pernambuco,
Department of Biochemistry, Recife-PE,
Brazil
<http://lattes.cnpq.br/7863845087003953>

Bruno Oliveira de Veras

Federal University of Pernambuco,
Department of Biochemistry, Recife-PE,
Brazil
<http://lattes.cnpq.br/7515860243444988>

Fernanda Miguel de Andrade

Federal University of Pernambuco,
Department of Biochemistry, Recife-PE,
Brazil
Sertão School of Medicine – FMS,
Arcoverde-PE, Brazil
<http://lattes.cnpq.br/8915203118340030>

Abstract: Plants have been manipulated for therapeutic purposes with the aim of preventing, treating and curing pathologies. Inhibition of cholinesterases contributes to increased bioavailability of the neurotransmitter acetylcholine in the synaptic cleft. The therapy for Alzheimer's disease (AD) consists of inhibiting cholinesterases, and natural products are rich sources of these inhibitors. This study aimed to evaluate the chemical composition and in vitro anticholinesterase activity of essential oils from *Hymenaea cangaceira* (OEHC) and *Eugenia gracilima* (OEEG). The essential oils were obtained by hydrodistillation and evaluated for composition in a gas chromatograph, coupled to a mass spectrometer (GC-MS) and flame ionization detector (GC-DIC). Cholinesterase inhibition was performed in in vitro colorimetric assays, using acetylcholinesterase (AChE) and butyrylcholinesterase (BChE), the results being expressed in $\mu\text{g/mL}$ of the concentration required for IC_{50} . Neostigmine was used as a reference standard. The OEHC presented 15 compounds and the OEEG 42 compounds. In the evaluation of AChE inhibition, the OEHC presented an IC_{50} of $0.015\mu\text{g/mL}$, being statistically lower than the OEEG (IC_{50} of $18.30\mu\text{g/mL}$). The OEHC against the BChE enzyme presented an IC_{50} of $0.041\mu\text{g/mL}$, being lower than the OEEG, which obtained an IC_{50} of $8.67\mu\text{g/mL}$. OEHC obtained greater inhibitory activity on AChE and BChE when compared to OEEG. The results suggest that OEHC has potential for use in the clinical treatment of neurodegenerative diseases, since it decreases AChE and BChE activity in vitro.

Keywords: Alzheimer's disease. Acetylcholinesterase. Butyrylcholinesterase. Enzyme Inhibition. Natural products.

INTRODUCTION

Plants have been manipulated for therapeutic purposes since antiquity with the aim of preventing, treating and curing many pathologies (DE CARVALHO et al., 2013). The progress of science has strengthened research on medicinal plants, associating their chemical composition with their effects, often attesting to their popular use (CAVALCANTE et al., 2013).

Natural products have attracted attention in major research centers around the world. Brazil has a wide biological variety and has numerous plant species with potential commercialization in the pharmaceutical, food and cosmetic sectors, among others (DA SILVA et al., 2013). Among plant species that are part of the biological diversity of Brazil are *Hymenaea cangaceira* and *Eugenia gracillima*.

Hymenaea cangaceira belongs to the Fabaceae family. It is popularly known as jatobá-do-cangaço/cangaceiro. It was recently described by Pinto (2017) and is found in the Caatinga region of Northeast Brazil, more specifically in the states of Bahia, Ceará and Pernambuco. The species of *Eugenia gracillima* belongs to the Myrtaceae family. Its popular name is described as myrtle/guajuraia/pitanga-guabiroba. It is a native fruit tree in Chapada do Araripe, located in the state of Pernambuco (DE ARAÚJO et al., 2016). There are still no studies related to the two species regarding anticholinesterase activity.

Acetylcholinesterase (AChE) and Butyrylcholinesterase (BChE) are cholinesterase enzymes of the nervous system that catalyze the hydrolysis of acetylcholine (ACh) to acetic acid and choline in the synaptic cleft. The action of these enzymes reduces the levels of the neurotransmitter ACh and, consequently, the loss of communication between nerve cells, causing a decrease in brain function. Both enzymes have ACh

as a substrate, however, BChE has a lower affinity for the substrate compared to AChE (COLOVIC et al., 2013; ZILBEYAZ et al., 2018).

The reduction of the neurotransmitters ACh, noradrenaline, serotonin, somatostatin and glutamate are associated with several diseases linked to the degeneration of cholinergic neurons, including Alzheimer's disease (AD). AD is considered a public health problem by the World Health Organization (WHO), as it is the most common reason for dementia, accounting for approximately 60% to 80% of cases, affecting thousands of elderly people worldwide. AD, which is a disturbing neurocognitive disorder recognized by a gradual reduction of cognitive functions and memory, leads patients to emotional stress, behavior and personality changes, and can reduce independent daily life (SIBERSKI, 2012; MAIER; BARNIKOL, 2014); BONO, 2014; WHO, 2019).

Currently, therapy for AD consists of inhibiting AChE, as patients with this disease have low levels of ACh (FERRO; GUIMARÃES; GONÇALVES, 2019). AChE inhibitors, called anticholinesterases, increase the bioavailability of the neurotransmitter ACh in the synaptic cleft. This way, the inhibition of this enzyme contributes to the reestablishment of cell communication (ANAND; SINGH, 2013).

The importance of finding new therapeutic options for AD has motivated research aimed at evaluating in vitro and in vivo anticholinesterase activity of products obtained synthetically and/or naturally (natural products) (SILVA, 2018). Among the compounds that have shown enormous potential for application for the development of new drugs and, consecutively, the treatment of neurodegenerative diseases such as AD, are medicinal plants (NEVES et al., 2017).

Products obtained from medicinal plants, such as essential oils, have been reported as one of the innovative strategies for the development of new drugs, called herbal medicines, gaining considerable importance and visibility in recent decades (LOIZZO et al., 2009; JEON et al, 2011; MAJLESSI et al., 2012; CHAIYANA; OKONOGI, 2012). Based on this problem, the objective was to evaluate the chemical composition and in vitro anticholinesterase activity of the essential oils of *Hymenaea cangaceira* and *Eugenia gracillima*.

METHODOLOGY

PLANT MATERIAL

The leaves of *Hymenaea cangaceira* Pinto, Mansano & Azevedo and *Eugenia gracillima* Kiaersk were collected in April 2018, in Catimbau National Park, located in the municipality of Buíque, in the state of Pernambuco.

The botanical identification of the species was performed using exsiccates. To prepare the exsiccates, the collected sample was pressed and dried (37° C for 72 h) (ORLANDA, 2011). Then, the exsiccates were sent to the Herbário Dárdano de Andrade Lima, of the Instituto Agronômico de Pernambuco (IPA), being deposited under the numbers 84888 and 78999, respectively.

EXTRACTION OF ESSENTIAL OILS

The leaves were submitted to mechanical separation for disposal of materials attacked by pests or diseases. Then, they were washed and crushed in a knife mill, and the material was later transferred to a hydrodistillation flask with water in a 1:10 proportion (m/v), and then heated for 5 hours at 100 °C.

Essential oils were obtained by condensing water vapor in the condenser together with volatile components. To remove the water, the drying agent sodium sulfate anhydrous

was added to the mixture of essential oil and water. (Na_2SO_4). The oils were kept refrigerated at -4 °C in an amber bottle (DE VERAS et al., 2019).

ANALYSIS OF THE CHEMICAL COMPOSITION OF ESSENTIAL OILS

Essential oils were analyzed on a 5975C triple gas chromatograph (Palo Alto, CA, USA) equipped with an Agilent Technologies DB-5MS column (30 m x 0.25 mm x 0.25 μm). Aliquots of 1 mL in the 1:50 division of essential oils were injected in gas chromatography coupled to a mass spectrometer (CG-EM). Compounds were identified by comparing their mass spectra using MassFinder 4, NIST08 and Wiley Registry software.™ 9th Edition, integrated with Agilent MSD Productivity ChemStation software (Agilent Technologies, Palo Alto, USA) and the Retention Index. The samples were quantified by gas chromatography with a flame ionization detector. (CG-DIC), under the same conditions as the GC-MS in triplicate, to calculate the standard deviation of the percentage of peak area for each compound in the chromatogram (DE VERAS et al., 2019).

ASSESSMENT OF CHOLINESTERASE INHIBITORY ACTIVITY

Obtaining the enzymes

It was used AChE de *Electrophorus electricus* (Type VI-S) obtained from Sigma-Aldrich. The BChE used was extracted from the gills of *Crassostrea rhizophorae* mangrove oysters, by homogenization of 20 mg of tissue per mL of 0.5 mol/L Tris-HCl pH 8.0 and centrifugation at 10,000 rpm (rotations per minute) for 10 minutes. The supernatant (crude extract) was collected and kept at -20°C for further analysis (DE SOUZA et al., 2018).

Enzyme inhibition assay

The enzyme inhibition assays were performed by incubating 200 μL of 0.25 $\text{mmol}\cdot\text{L}^{-1}$ DTNB (Ellman's Reagent, dissolved in 0.5 $\text{mol}\cdot\text{L}^{-1}$ of Tris-HCl buffer pH 7.4), 10 μL of the enzymes (20 mg / mL of buffer), with 10 μL of essential oils (increasing concentrations from 0.01 to 1000 $\mu\text{g}/\text{mL}$) for 60 minutes, according to Assis et al. (2012). The tests were carried out at room temperature (25°C). The reaction started immediately after the addition of 62 mmol/L of S-acetylcholine or S-butyrylthiocholine (20 μL). Reactions were performed in quadruplicate in 96-well microplates. The plates were read by spectrophotometry at 405 nm at 0 and 180 s. As a blank, a 0.5 mol/L solution of Tris-HCl pH 8.0 was used instead of the essential oils. Neostigmine was used as a reference standard, at the same concentrations as the essential oils.

Results were expressed as percent residual activity, mean \pm SD (Standard Deviation), in contrast to inhibitor concentration in mg/mL . Neostigmine was used as the reference drug. Residual activity (%) was plotted against inhibitor concentration in $\mu\text{g}/\text{mL}$. Results were expressed as mean \pm SD of four replicates and statistically analyzed by nonlinear regression fitted to sigmoidal decay models using Microcal™ Origin® versão 8.0.

RESULTS AND DISCUSSION

ANALYSIS OF THE CHEMICAL COMPOSITION OF ESSENTIAL OILS

The analysis of the chemical composition of *Hymenaea cangaceira* essential oil (OEHC) indicated the presence of 15 chemical components, corresponding to 85.37% of the total components, all of which are classified as sesquiterpene hydrocarbons. Among the major components, α -Copaene (6.34% \pm 0.17), β -Elemene (7.05% \pm 0.12), (E)-Caryophyllene (23.38% \pm 0.58), α -Guaiene (9.75% \pm 0.07), α -

-Humulene (4.65% \pm 0.12) and Germacrene D (14.66% \pm 0.14) (Table 1).

De Veras et al. (2020) reported that OEHC has α -Copaene, β -Elemene, (E)-Caryophyllene, α -Guaiene, α -Humulene and Germacrene D as main constituents, being present at the same concentrations as in the present study.

Several activities have been described with several compounds found in OEHC. α -Copaene has antimicrobial, anti-inflammatory and healing activity (BRITO et al., 2005). β -Elemene is presented as anticancer, antihypertensive and anti-inflammatory (LI et al., 2009; YVON et al., 2012; LI et al., 2013; CAI et al., 2013; ZOU et al., 2013; ; ZHANG et al., 2013; DONG; YIN; WANG, 2013). (E)-Caryophyllene has antimicrobial and antifungal activity (GARG; SIDDIQUI, 1992; FORMISANO et al., 2006). α -Guaiene is described as an inhibitor of cyclooxygenase, 5-lipoxygenase and acetylcholinesterase (IBRAHIM et al., 2016). Fernandes et al. (2007) described anti-inflammatory activity for α -Humulene. Germacrene D has been reported to be a potent antimicrobial for gram-positive bacteria (DUARTE, 2006).

A study carried out with essential oils of *Piper anonifolium* and *Piper hispidum* showed anticholinesterase activity, stating that the inhibition was due to the presence of sesquiterpene compounds found in their oils (DA SILVA et al., 2014). Some of the components were also described in the present work as: α -Copaene, β -Elemene, α -Humulene and Germacrene D.

The analysis of the chemical composition of *Eugenia gracillima* essential oil (OEEG) showed the presence of 42 compounds, corresponding to more than 99% of the total components. OEEG is mainly composed of 90.11% of sesquiterpene hydrocarbons and 9.57% are composed of oxygenated

Peak	Compound	RI Cal	RI Lit	Percentage (%)	D.P.
1	δ -Elemene	1341	1335	2.21	0.06
2	α -Copaene	1380	1374	6.34	0.17
3	β -Elemene	1396	1389	7.05	0.12
4	Cyperene	1403	1398	2.68	0.06
5	(E)-Caryophyllene	1424	1417	23.38	0.58
6	γ -Elemene	1438	1434	2.18	0.13
7	α -Guaiene	1443	1437	9.75	0.07
8	α -Humulene	1458	1452	4.65	0.12
9	γ -Muuroolene	1481	1478	1.16	0.10
10	Germacrene D	1486	1480	14.66	0.14
11	δ -Selinene	1496	1492	1.14	0.04
12	Bicyclogermacrene	1501	1500	4.59	0.02
13	α -Bulnesene	1511	1509	2.84	0.04
14	δ -Cadinene	1529	1522	2.21	0.06
15	Germacrene B	1563	1559	0.53	0.02
Total				85.37	
Others				14.63	

RICal- Relative experimental retention indices and RILit - literature retention indices; DP. Standard deviation.

Table 1. Chemical composition of OEHC, highlighted in bold the main components.

sesquiterpenes. The most abundant constituents were δ -Elemene ($6.06\% \pm 0.08$), Bergamotene ($4.89\% \pm 0.06$), γ -Muuroolene ($15.60\% \pm 0.22$), Germacrene D ($16.10\% \pm 0.38$), Bicyclogermacrene (8, $53\% \pm 0.09$), δ -Cadinene ($6.23\% \pm 0.16$) and Germacrene B ($7.43\% \pm 0.62$) (Table 2).

Sampaio et al. (2019) reported that OEEg has δ -Elemene, γ -Muuroolene, Germacrene D, Bicyclogermacrene and Germacrene B as main constituents, being present at the same concentrations as in the present study.

Several species of *Eugenia* spp. have great biological potential such as antimicrobial, antifungal, antitumor, among others. These can be attributed to the volatile chemical composition present in its essential oils (MAGINA, 2008). δ -Elemene exerts

anticancer activity (LU et al., 2012). Bergamotene has antioxidant activity (DE MELO et al. 2011). γ -Muuroolene has antimicrobial activity (SILVA et al., 2010; CHAIBUB et al., 2013). Germacrene D has insecticidal, antibacterial and fungicidal action (ALMEIDA; DELACHIAVE; MARQUES, 2005). Bicyclogermacrene has a high potential for antimicrobial action (CONSTANTIN et al., 2001; CYSNE et al., 2005; SOUSA et al., 2010). δ -Cadinene has antitumor and antimicrobial activity (SKAL TSA et al., 2003; WRIGHT et al., 2007). Germacrene B has antifungal activity (COSTA et al., 2010).

Besides, in the study carried out by Da Silva et al. (2014), some OEEg compounds were also seen in the essential oils of *Piper*

Peak	Compound	RI Cal	RI Lit	Percentage (%)	D.P.
1	δ-Elemene	1336	1335	6.06	0.08
2	α -Cubebene	1345	1345	0.12	0.00
3	α -Ylangene	1376	1373	0.50	0.02
4	Isoledene	1375	1374	1.30	0.03
5	Copaene	1377	1374	0.60	0.00
6	Guaia-6,9-diene	1380	1379	0.68	0.04
7	β -Bourbonene	1387	1387	0.71	0.00
8	β -Cubebene	1388	1387	0.11	0.00
9	β -Elemene	1390	1389	1.41	0.08
10	Sativene	1392	1390	0.21	0.00
11	α -Gurjunene	1410	1409	4.03	0.02
12	β -Copaene	1432	1430	0.31	0.05
13	Bergamotene	1433	1432	4.89	0.06
14	γ -Elemene	1434	1434	0.73	0.00
15	α -Guaiene	1437	1437	1.80	0.08
16	Cadina-3,5-diene	1449	1448	0.26	0.09
17	α -Humulene	1454	1452	1.64	0.11
18	Alloaromadendrene	1458	1458	0.90	0.00
19	Caryophyllene-9-epi-(e)	1461	1464	0.40	0.00
20	Cumacrene	1470	1470	0.27	0.00
21	γ-Muurolene	1478	1478	15.60	0.22
22	γ -Amorphene	1482	1483	3.14	0.00
23	Germacrene D	1483	1484	16.10	0.38
24	β -Selinene	1490	1489	0.16	0.00
25	Bicyclogermacrene	1501	1500	8.53	0.09
26	α -Muurolene	1503	1500	0.73	0.00
27	β -Guaiene	1504	1502	0.84	0.01
28	δ-Cadinene	1524	1522	6.23	0.16
29	Zonarene	1528	1528	0.93	0.00
30	α -Cadinene	1537	1537	2.33	0.17
31	Selina-3,7(11)-diene	1548	1545	0.15	0.00

32	Elemol	1549	1548	0.61	0.00
33	Germacrene B	1561	1559	7.43	0.62
34	Spathulenol	1578	1577	0.67	0.00
35	Thujopsan-2 β -ol	1589	1588	0.29	0.00
36	Globulol	1591	1590	1.03	0.05
37	1h-cycloprope-azulen-4-ol	1592	1592	1.01	0.10
38	Viridiflorol	1593	1592	0.42	0.00
39	Isospathulenol	1626	1625	1.25	0.09
40	Torreyol	1644	1644	1.27	0.12
41	Cubenol	1645	1645	3.58	0.19
42	α -Cadinol	1656	1654	0.45	0.00
Total				99.68	
Outros				0.32	

RICal- Relative experimental retention indices and RILit - literature retention indices; DP. Standard deviation.

Table 2. Chemical composition of OEEg, the main components highlighted in bold.

Sample	Inhibition of the activity of AChE IC ₅₀ (μ g/mL)	Inhibition of the activity of BChE IC ₅₀ (μ g/mL)
OEHC	0.015 \pm 0.000 ^{3a}	0.041 \pm 0.005 ^a
OEEg	18.30 \pm 0.0041 ^c	8.67 \pm 0.0012 ^c
Neostigmine	0.16 \pm 0.0 ^{1b}	1.9 \pm 0.0006 ^b

Data represent the mean \pm SD. ^{abc} Means followed by different letters in the column are significantly different (p < 0,05) by the test of Tukey.

Table 3. AChE and BChE inhibitory activity by OEHC and OEEg.

anonifolium and *Piper hispidum* that also have anticholinesterase activity, such as: δ -Elemene, γ -Muuroleone, Germacrene D and Bicyclgermacrene. The chemical compositions found need to be isolated to understand whether the anticholinesterase activity acted synergistically, or is performed by some specific compounds alone. Because the biological actions of essential oils are not exactly associated with the amount of the main components, but can also be related to interactions between the largest and smallest compositions, which generate synergies and/or antagonistic responses (AAZZA; LYOUSSI; MIGUEL, 2011; TAK; JOVEL; ISMAN, 2016).

ENZYME INHIBITION

For the enzyme inhibition assay, the inhibitory concentration IC₅₀ was used. IC₅₀ values represent the average inhibition of a compound (50% inhibition), that is, it corresponds to the dose capable of inhibiting 50% of the enzyme. Thus, the lower the IC₅₀, the more active a compound is.

The OEHC showed the highest inhibition of AChE statistically, presenting an IC₅₀ value of $0.015 \pm 0.0003 \mu\text{g/mL}$. OEEg, despite having shown inhibition of AChE (IC₅₀ $18.30 \pm 0.0041 \mu\text{g/mL}$), had lower activity compared to OEHC. The drug Neostigmine, used as a standard, had an IC₅₀ of $0.16 \pm 0.01 \mu\text{g/mL}$ (Table 3).

In the evaluation of the inhibitory activity against the BChE enzyme, it was observed that OEHC resulted in a high inhibitory activity with an IC₅₀ of $0.041 \pm 0.005 \mu\text{g/mL}$. The Neostigmine standard presented an IC₅₀ of $1.9 \pm 0.0006 \mu\text{g/mL}$. The OEEg obtained inhibitory activity with an IC₅₀ of $8.67 \pm 0.0012 \mu\text{g/mL}$, the inhibitory action being not as significant as compared to the OEHC (Table 3).

A study already carried out with *Hymenaea cangaceira* relatou que o óleo essencial

da planta possui potencial antioxidante, atividade antimicrobiana e atividade antinociceptiva (DE VERAS et al., 2020).

Regarding anticholinesterase activity, this is the first study carried out in OEHC and OEEg. Several studies with essential oils have demonstrated their potential to inhibit key enzymes in the control of diseases such as AD, such as the essential oil of *Citrus sinensis*, which had an AChE enzyme inhibitory activity of IC₅₀ = $63 \mu\text{g/mL}$ (SÁ et al., 2012). A study carried out with the species of *Salvia lavandulaefolia* showed inhibitory activity against the AChE enzyme in the value of IC₅₀ = $50 \mu\text{g/mL}$ (SAVELEV et al., 2003). Another research, with the essential oils of *Eucalyptus camaldulensis* and *Ocimum canum*, showed inhibition of AChE with IC₅₀ value = $18.98 \mu\text{g/mL}$ and IC₅₀ = $36.16 \mu\text{g/mL}$, respectively (KIENDREBEOGO et al., 2011). The OEHC and the OEEg in the present study showed superior anticholinesterase activity, in relation to the essential oils of other species already mentioned, confirming their effectiveness and better inhibition of the enzyme, revealing the importance of researching new therapies for AD.

CONCLUSION

OEHC obtained greater inhibitory activity on AChE and BChE enzymes compared to OEEg. The results suggest that OEHC has potential for use in the clinical treatment of neurodegenerative diseases, such as AD, since it significantly inhibited AChE and BChE activity in vitro. Therefore, these results are seen as innovative strategies for the development of new drugs/herbal medicines.

REFERENCES

- AAZZA, S.; LYOUSSI, B.; MIGUEL, M.G. Antioxidant and antiacetylcholinesterase activities of some commercial essential oils and their major compounds. **Molecules**, v. 16, n. 9, p. 7672–7690, sep. 2011.
- ALMEIDA, L. F. R.; DELACHIAVE, M. E. A.; MARQUES, M. O. M. Composição do óleo essencial de rubim (*Leonurus sibiricus* L. - Lamiaceae). **Revista Brasileira de Plantas Mediciniais**, Botucatu, v. 8, n. 1, p. 35-38, 2005.
- ANAND, P.; SINGH, B. A review on cholinesterase inhibitors for alzheimer's disease. **Archives of Pharmacal Research**, Patiala, v. 36, n. 4, p. 375–399, apr. 2013.
- ASSIS, C. R. D. et al. Comparative effect of pesticides on brain acetylcholinesterase in tropical fish. **Science of the Total Environment**, Recife, v. 441, p. 14-150, dec. 2012.
- BONO, G. F. **Investigação de associação entre a butirilcolinesterase e a doença de alzheimer**. 2014. 106f. Dissertação (Mestrado) - Universidade Federal do Paraná, Curitiba.
- BRITO, M. V. H. et al. Efeito do óleo de copaíba nos níveis séricos de ureia e creatinina em ratos submetidos à síndrome de isquemia e reperfusão renal. **Acta Cirúrgica Brasileira**, São Paulo, v. 20, n. 3, p. 243-246, mai./jun. 2005.
- CAI DY. et al. Synergistic effect of beta-elemene injection combined paclitaxel injection on human breast cancer MB-468 cells: an *in vitro* study. **Zhongguo Zhong Xi Yi Jie He Za Zhi**, China, v. 33, n. 7, p. 978-982, jul. 2013.
- CAVALCANTE, G. M. et al. Atividade antimicrobiana de *Artocarpus heterophyllus* Lam. (Moraceae) sobre o desenvolvimento de *Streptococcus pneumoniae* e *Escherichia coli*. **Scientia Plena**, v. 9, n. 2, p. 1-7. 2013.
- CHAIBUB, B. A. et al. Composição química do óleo essencial e avaliação da atividade antimicrobiana do óleo essencial, extrato etanólico bruto e frações das folhas de *Spiranthera odoratissima* A St.-Hil. **Revista Brasileira de Plantas Mediciniais**, Botucatu, v. 15, n. 2, p. 225-229, 2013.
- CHAIYANA, W.; OKONOJI, S. Inhibition of cholinesterase by essential oil from food plant. **Phytomedicine**, Thailand, v. 19, n. 8-9, p. 836-839, apr. 2012.
- COLOVIC, M. B. et al. Acetylcholinesterase inhibitors: pharmacology and toxicology. **Current neuropharmacology**, Belgrade, v. 11, n. 3, p. 315-335, may. 2013.
- CONSTANTIN, M. B. et al. Essential oils from *Piper cernuum* and *Piper regnellii*: antimicrobial activities and analysis by CG/MS and C-NMR. **Planta Med**, São Paulo, v. 67, n. 8, p. 771-733, apr. 2001.
- COSTA, D. P. et al. Influence of fruit biotypes on the chemical composition and antifungal activity of the essential oils of *Eugenia uniflora* leaves. **Journal of the Brazilian Chemical Society**, São Paulo, v. 21, n. 5, p. 851-858, feb. 2010.
- CYSNE J. B. et al. Leaf essential oils of four *Piper* species from the State of Ceará - Northeast of Brazil. **Journal of the Brazilian Chemical Society**, São Paulo, v. 16, n. 6B, p. 1378-1381, nov./dec. 2005.
- DA SILVA, J. K. R. et al. Essential oils of Amazon *Piper* species and their cytotoxic, antifungal, antioxidant and anti-cholinesterase activities. **Industrial Crops and Products**, v. 58, p. 55–60, jul.2014.
- DA SILVA, S. R. F. **Farmacocinética do diazepam**. 2013. 79f. Monografia (Mestrado) - Universidade Fernando Pessoa, Porto.
- DE ARAÚJO, D. R. et al. Characterization of ripening stages of myrtle fruit. **Revista Brasileira de Fruticultura**, Jaboticabal, v. 38, n. 2, p. 1-10, mar/apr. 2016.
- DE CARVALHO, R. B. F. et al. Composição química e atividade anticolinesterásica de uma fração ativa do extrato de folhas de *Citrus limon* (L.) Burm. **Química Nova**, v. 36, n. 9, p. 1375-1379, ago. 2013.
- DE MELO, N. I. et al. Schistosomicidal activity of the essential oil of *Ageratum conyzoides* L. (Asteraceae) against adult *Schistosoma mansoni* worms. **Molecules**, v. 16, n. 1, p. 762-773, 2011.
- DE SOUZA, P. R. et al. Acetylcholinesterase of mangrove oyster *Crassostrea rhizophorae*: A highly thermostable enzyme with promising features for estuarine biomonitoring. **Aquatic Toxicology**, v. 197, p. 109–121, 2018.

- DE VERAS, B. O. et al. Chemical composition and evaluation of the antinociceptive, antioxidante and antimicrobial effects of essential oil from *Hymenaea cangaceira* (Pinto, Mansano & Azevedo) native to Brazil: A natural medicine. **Journal of Ethnopharmacology**, v. 247, jan. 2020.
- DE VERAS, B. O. et al. *Algrizea Minor* Sobral, Faria & Proença (Myrteae, Myrtaceae): chemical composition, antinociceptive, antimicrobial and antioxidant activity of essential oil. **Natural Product Research**, v. 23, p. 1-5, apr. 2019.
- DONG Q.; YIN T.; WANG G. Potential applications of beta-elemene in anti-atherosclerosis and anti-restenosis. **Sheng Wu Yi Xue Gong Cheng Xue Za Zhi**, China, v. 30, n. 3, p. 656-660, jun. 2013.
- DUARTE, M.C.T. Atividade antimicrobiana de plantas medicinais e aromáticas utilizadas no Brasil. **Revista Multiciência**, Unicamp, n. 7, p. 1-16. 2006.
- FERNANDES, E. S. et al. Anti-inflammatory effects of compounds alpha-humulene and (-)-trans-caryophyllene isolated from the essential oil of *Cordia verbenacea*. **European Journal Of Pharmacology**, v. 569, n. 3, p. 228-236, may. 2007.
- FERRO, E. D. S.; GUIMARÃES, A. P.; GONÇALVES, A. S. L. Estudo de ancoramento molecular (docking) de potenciais inibidores da acetilcolinesterase humana (*HssAChE*). **Anais do XIII Congresso Brasileiro de Engenharia Química em Iniciação Científica**, São Paulo, v. 1, n. 6, p. 2639-2645, ago. 2019.
- FORMISANO, C. et al. Chemical composition and antimicrobial activity of the essential oil of *Phlomis ferruginea* Ten. (Lamiaceae) growing wild in Southern Italy. **Flavour and Fragrance Journal**, v. 21, n. 5, p. 848-851, sep./oct. 2006.
- GARG, S. C.; SIDDIQUI, N.; Antifungal activity of some essential oil isolates. **Pharmazie**, v. 47, n. 6, p. 467-468, jun. 1992.
- IBRAHIM, M. S. et al. Cyclooxygenase, 5-lipoxygenase and acetylcholinesterase inhibitory effects of fractions containing α -guaiene and oil isolated from the root of *Xylocarpus moluccensis*. **Research Journal of Medicinal Plants**, v. 10, n. 4, p. 286-294, apr. 2016.
- JEON, S. et al. SuHeXiang Wan Essential Oil Alleviates Amyloid Beta Induced Memory Impairment Through Inhibition of Tau Protein Phosphorylation in Mice. **The American Journal of Chinese Medicine**, v. 39, n. 5, p. 917-932. 2011.
- KIENDREBEOGO, M. et al. Antiacetylcholinesterase and antioxidant activity of essential oils from six medicinal plants from Burkina Faso. **Brazilian Journal of Pharmacognosy**, v. 21, n. 1, p. 63-69, jan./feb. 2011.
- LI, Q. Q. et al. β -Elemene enhances susceptibility to cisplatin in resistant ovarian carcinoma cells via downregulation of ERCC-1 and XIAP and inactivation of JNK. **International Journal Of Oncology**, v. 43, n. 3, p. 721-728, sep. 2013.
- LI, Q. Q. et al. β -Elemene, a novel plant-derived antineoplastic agent, increases cisplatin chemosensitivity of lung tumor cells by triggering apoptosis. **Oncology Reports**, v. 22, n. 1, p. 161-170, jul. 2009.
- LOIZZO, M. R. et al. *In vitro* biological activity of *Salvia leriifolia* Benth essential oil relevant to the treatment of Alzheimer's disease. **Journal of Oleo Science**. v. 58, p. 443-446, 2009.
- LU, J-J. et al. Anti-cancer properties of terpenoids isolated from *Rhizoma Curcumae* - A review. **Journal of Ethnopharmacology**, v. 143, n. 2, p. 406-411, sep. 2012.
- MAGINA, M. D. A. **Estudo fitoquímico e biológico de espécies do gênero *Eugenia***. 2008. 178f. Tese (doutorado) - Universidade Federal de Santa Catarina, Centro de Ciências Físicas e Matemáticas. Programa de Pós-Graduação em Química, Florianópolis.
- MAIER, W.; BARNIKOL, U. B. Neurokognitive Störungen im DSM-5: Durchgreifende Änderungen in der Demenzdiagnostik. **Der Nervenarzt**, v. 85, n. 5, p. 564-570, 2014.
- MAJLESSI, N. et al. Amelioration of amyloid β -induced cognitive deficits by *Zataria multiflora* Boiss. essential oil in a rat model of Alzheimer's disease. **CNS Neuroscience & Therapeutics**, v. 18, n. 4, p. 295-301, apr. 2012.
- NEVES, A. M. et al. Caracterização Química e o Potencial Antimicrobiano de Espécies do Gênero *Senna* Mill. (Fabaceae). **Revista Virtual de Química**, v. 9, n. 6, 2017.

- ORLANDA, J. F. F. **Estudo da composição química e atividade biológica do óleo essencial de *Ruta graveolens* Linneau (RUTACEAE)**. 2011. 105p. Tese (Doutorado) - Universidade Federal da Paraíba, Centro de Ciências Exatas e da Natureza. Programa de Pós-Graduação em Química, João Pessoa.
- PINTO, R. B. **Systematic studies in the *Hymenaea* clade and a taxonomic revision of the genus *Hymenaea* L. (Leguminosae, Detarioideae)**. 2017. 203f. Tese (Doutorado em Biologia Vegetal) - Universidade Estadual de Campinas, Campinas.
- SÁ, C. G. et al. Efeito do tratamento agudo do óleo essencial de *Citrus sinensis* (L) Osbeck na aquisição da memória espacial de ratos avaliada no labirinto aquático de Morris. **Revista de Ciências Farmacêuticas Básica e Aplicada**, v. 33, n. 2, p. 211-215, 2012.
- SAMPAIO, M. G. V. et al. Chemical composition, antioxidant and antiprotozoal activity of *Eugenia gracillima* Kiaersk. leaves essential oil. **Natural Product Research**, p. 1-5, 2019.
- SAVELEV, S. et al. Synergistic and antagonistic interactions of anticholinesterase terpenoids in *Salvia lavandulaefolia* essential oil. **Pharmacology, Biochemistry and Behavior**, v. 75, n. 3, p. 661-668, 2003.
- SIBERSKI, J.; MS.; CMC. Dementia and DSM-5: Changes, cost, and confusion. **Aging Well**, v. 5, n. 6, p. 12, 2012. Disponível em: <http://www.todaygeriatricmedicine.com/archive/110612p12.shtml>. Acesso em: 07/09/2019.
- SILVA, C. D. S. et al. Avaliação *in vitro* do efeito de frações de *O. glaziovii* frente às enzimas acetilcolinesterase e butirilcolinesterase. **Anais Seminário de Iniciação Científica**. Feira de Santana, n. 22, p. 1, 2018.
- SILVA, L. L. et al. Composição química, atividade antibacteriana *in vitro* e toxicidade em *Artemia salina* do óleo essencial das inflorescências de *Ocimum gratissimum* L., Lamiaceae. **Revista Brasileira de Farmacognosia**, Curitiba, v. 20, n. 5, p. 700-705, 2010.
- SKAL TSA, H. D. et al. Essential oil analysis and antimicrobial activity of eight *Stachys* species from Greece. **Phytochemistry**, v. 64, n. 3, p. 743-752, 2003.
- SOUZA, E. O. et al. Chemical composition and resistance-modifying effect of the essential oil of *Lantana camara* Linn. **Pharmacognosy Magazine**, v. 6, n. 22, p. 7-82, apr. 2010.
- TAK, J.-H.; JOVEL, E.; ISMAN, M. B. Comparative and Synergistic Activity of *Rosmarinus officinalis* L. Essential Oil Constituents Against the Larvae and an Ovarian Cell Line of the Cabbage Looper, *Trichoplusia ni* (Lepidoptera: Noctuidae). **Pest Management Science**, v. 72, n. 3, p. 474-480, mar. 2016.
- WORLD HEALTH ORGANIZATION (WHO). 2019. **Dementia**. Disponível em: <http://www.who.int/mediacentre/factsheets/fs362/en/>. Acesso em: 07/09/2019.
- WRIGHT, B. S. et al. Cytotoxic leaf essential oils from Neotropical Lauraceae: Synergistic effects of essential oil components. **Natural Product Communications**, v. 2, n. 12, p. 1241-1244, 2007.
- YVON, Y. et al. Relation between chemical composition or antioxidant activity and antihypertensive activity for six essential oils. **Journal of Food Science**, v. 77, n. 8, p. 184-191, aug. 2012.
- ZHANG, Y. et al. The Role of E3 Ubiquitin Ligase Cbl Proteins in β -Elemene Reversing Multi-Drug Resistance of Human Gastric Adenocarcinoma Cells. **International Journal of Molecular Sciences**, v. 14, n. 5, p. 10075-10089, may. 2013.
- ZILBEYAZ, K. et al. Effects of aryl methanesulfonate derivatives on acetylcholinesterase and butyrylcholinesterase. **Journal of Biochemical and Molecular Toxicology**, v. 32, n. 11, p. 1-4. 2018.
- ZOU, B. et al. β -Elemene and taxanes synergistically induce cytotoxicity and inhibit proliferation in ovarian cancer and other tumor cells. **Anticancer Research**, v. 33, n. 3, p. 929-940, mar. 2013.