

IMPACT OF PRENATAL CARE ON OBSTETRIC OUTCOMES: A NARRATIVE REVIEW OF THE LITERATURE

Andressa Aguiar da Silva

<https://lattes.cnpq.br/8911547342317913>

Mariana Dantas Moreira

<https://lattes.cnpq.br/5874478828844366>

Ivan Lucas Bonizioli Barcellos

<https://orcid.org/0009-0000-9478-7008>

Yolaida Betancourt Casanova

<https://lattes.cnpq.br/3763322176929828>

Deilton Sousa Costa

<https://lattes.cnpq.br/1351701225557512>

Fernanda Ferreira Gouveia

<https://lattes.cnpq.br/2875545470504486>

Eduardo Soares Marques Guimarães

<https://orcid.org/0009-0008-4978-1262>

Bianca Yumi Takano

<https://lattes.cnpq.br/418956119270165489>

Aline Santos de Almeida

<https://lattes.cnpq.br/5945758395564308>

Ketyla Albino Linhares

<https://lattes.cnpq.br/5414248369063517>

Joyce Almeida Né da Silva

<https://orcid.org/0000-0002-5125-0048>

Janaina Bojikian da Costa Vital Juliatto

<https://lattes.cnpq.br/2596242858222029>

All content in this magazine is licensed under a Creative Commons Attribution License. Attribution-Non-Commercial-Non-Derivatives 4.0 International (CC BY-NC-ND 4.0).



Abstract: Objective: To analyze the impact of prenatal care on obstetric outcomes. Literature Review: Prenatal care, through preventive actions, seeks to ensure the healthy development of pregnancy and enable the birth of a healthy baby, while preserving the health of the baby and the mother. Adequate prenatal care requires at least six consultations during pregnancy (Ministry of Health, 2006b). If the pregnancy is not classified as high-risk, at least one consultation is recommended in the first trimester of pregnancy, two in the second and three in the third. Pregnant women must be seen until they reach labor or reach the risk period for post-maturity, around the 42nd week. According to the Ministry of Health (2006b), prenatal care is concluded only on the 42nd day after delivery, when the postpartum consultation is held. These determinations may vary, depending on the risks presented by the pregnant woman, and in all cases adherence to prenatal care is essential (Buchabqui, Abeche & Brietzke, 2001). Final considerations: It is important to systematically evaluate prenatal care for pregnant women. This can be done by creating a space to listen to these pregnant women and also to the professionals involved in the care. This information is very important for improving prenatal care, understanding the motivations of pregnant women who do not adhere to it, and providing quality prenatal care in medical and psychological terms to all pregnant women.

Keywords: pregnant women, prenatal, neonatal.

INTRODUCTION

Pregnancy is a very important experience in the life of a woman and her family. During pregnancy, some physiological changes occur that involve all organic systems, generating expectation, emotion, anguish, concern and discoveries. Therefore, it is necessary to be aware of all these changes in order to provide adequate support for the health of the pregnant woman. Complicated pregnancies, whose particular characteristics can pose risks to the mother or fetus, are called “high risk”.

Therefore, prenatal care and comprehensive care for women’s health must be structured to meet the real needs of the pregnant population. Health care actions must be focused on the target audience of the area covered by the basic health unit (UBS), ensuring continuity of care, guidance and verification of these actions on perinatal and maternal health. In this context, prenatal support integrates precaution, behaviors and attitudes in favor of the pregnant woman.

This care occurs from the beginning of pregnancy until labor, and also aims to recognize, treat or verify the existence of pathologies; avoid complications during pregnancy and childbirth; provide pleasant maternal health and good fetal development; and reduce fetal and maternal morbidity and mortality rates. In 2000, the MS began implementing the Humanization Program in Prenatal and Birth Care (PHPN), through Ordinance GM No. 569, of June 1, 2000, with the objective of promoting actions aimed at reducing maternal morbidity and mortality and improving perinatal results, with the strategy of humanizing care.

BIBLIOGRAPHICAL REVIEW

Prenatal care, through preventive measures, seeks to ensure the healthy development of pregnancy and enable the birth of a healthy baby, while preserving the health of the baby and the mother. Studies have shown that qualified prenatal care is associated with a reduction in negative perinatal outcomes, such as low birth weight and prematurity, in addition to reducing the chances of obstetric complications, such as eclampsia, gestational diabetes and maternal deaths.

All guidance provided by health professionals to pregnant women during prenatal care is an important part of this care process. Although relevant, however, national studies have identified failure by health professionals to provide guidance on pregnancy, the importance of and techniques for breastfeeding, how to prepare for childbirth and basic care for the newborn.

In this sense, Primary Health Care (PHC) is a strategic space for low-risk, high-quality prenatal care. In Brazil, PHC, guided by the National Primary Care Policy (PNAB), emphasizes that it is the responsibility of the health team to welcome and provide health care to pregnant women and children, including disease prevention, health promotion, and treatment of problems that occur during pregnancy through the postpartum period, as well as child care. In this scenario, shared work among health professionals allows for different perspectives on prenatal care practices, ensuring comprehensive care and increasing the potential for resolution.

Adequate prenatal care requires at least six consultations during pregnancy (Ministry of Health, 2006b). If the pregnancy is not classified as high-risk, at least one consultation is recommended in the first trimester of pregnancy, two in the second and three in the third. Pregnant women must be seen until they reach labor or reach the risk period for post-

maturity, around the 42nd week. According to the Ministry of Health (2006b), prenatal care is only completed on the 42nd day after delivery, when the postpartum consultation is held. These determinations may vary, depending on the risks presented by the pregnant woman, and in all cases adherence to prenatal care is essential (Buchabqui, Abeche & Brietzke, 2001). It is worth noting that these recommendations are basically medical and do not mention the importance of assessing and meeting the emotional demands of the pregnant woman.

Such demands are, in some cases, at least partially addressed in support groups for pregnant women held in health centers, which have proven to be positive, although they are not a universal procedure (Falcone, Mäder, Nascimento, Santos & Nóbrega, 2005; Souza & Carvalho, 2003).

In addition to the factors associated with the quality of care, socioeconomic status has been identified as a factor that makes it difficult for Brazilian pregnant women to attend prenatal consultations, due to the difficulties in access that this vulnerability can cause (Silveira, Santos & Costa, 2001). Added to this are the long lines, long waiting times for consultations, the insensitive way in which some professionals deal with their patients, the lack of family support, difficulties with transportation, and cultural and language differences (Cook et al., 1999). The authors also mention that personal problems and psychological distress, such as depression, anxiety, stress or ambivalence regarding pregnancy, can influence adherence to prenatal care (Cook et al., 1999; Sable & Wilkinson, 1999).

Therefore, prenatal care appears to be an important time to offer emotional and social support to women. The mother-baby bond is certainly influenced by external and contextual factors in the pregnant woman's

life, one of which is social support (Huth-Bocks, Levendosky, Bogat & Von Eye, 2004). The authors state that mothers need to create and maintain a support network so that they can achieve the goals of motherhood. Corroborating these findings, Melender and Lauri (2002) identified that problems reported by North American mothers during pregnancy were associated with social support and experiences related to prenatal health care.

FINAL CONSIDERATIONS

It is important to systematically evaluate prenatal care for pregnant women. This can be done by creating a space to listen to these pregnant women and also to the professionals involved in providing care. This information is very important for improving prenatal care, understanding the motivations of pregnant women who do not adhere to it, and providing quality prenatal care in medical and psychological terms to all pregnant women. Understanding this period is not only extremely important for the immediate health issues of the pregnant woman and the baby, but also for motherhood, fatherhood and the baby's development.

REFERENCES

- PICCININI, Cesar Augusto et al. Percepções e sentimentos de gestantes sobre o pré-natal. *Psicologia: Teoria e Pesquisa*, v. 28, p. 27-33, 2012.
- CARNEIRO, Ana Beatriz Farias et al. A importância do pré-natal na prevenção de complicações durante a gestação. *Revista Brasileira Interdisciplinar de Saúde-ReBIS*, v. 4, n. 4, 2022.
- DA SILVA BOMFIM, Vitoria Vilas Boas et al. A importância do pré-natal no diagnóstico e tratamento da sífilis congênita. *Revista Eletrônica Acervo Saúde*, v. 13, n. 7, p. e7969-e7969, 2021.
- MARTINS, Quitéria Pricila Mesquita et al. Conhecimentos de Gestantes no pré-natal: Evidências para o cuidado de Enfermagem. *SANARE-Revista de Políticas Públicas*, v. 14, n. 2, 2015.
- MARQUES, Bruna Leticia et al. Orientações às gestantes no pré-natal: a importância do cuidado compartilhado na atenção primária em saúde. *Escola Anna Nery*, v. 25, n. 1, p. e20200098, 2020.
- CARDOSO, Soraya Lopes et al. Ações de promoção para saúde da gestante com ênfase no pré-natal. *Revista Interfaces: Saúde, Humanas e Tecnologia*, v. 7, n. 1, p. 180-186, 2019.
- DIAS, Ernandes Gonçalves et al. Percepção das gestantes quanto à importância das ações educativas promovida pelo enfermeiro no pré-natal em uma unidade básica de saúde. *Revista Gestão & Saúde*, v. 6, n. 3, p. ág. 2695-2710, 2015.
- BARRETO, Camila Nunes et al. Atenção pré-natal na voz das gestantes. *Revista de Enfermagem UFPE on line*, v. 7, n. 6, p. 4354-4363, 2013.
- BRITO, Lucas de Moraes Escorcio et al. A importância do pré-natal na saúde básica: uma revisão bibliográfica. *Research, Society and Development*, v. 10, n. 15, p. e51101522471-e51101522471, 2021.
- CARVALHO, Silas Santos; DE OLIVEIRA, Ludmila Freitas. Percepção de adolescentes gestantes sobre a assistência de enfermagem ao pré-natal. *Enfermagem em Foco*, v. 11, n. 3, 2020.
- DA SILVA BARROS, Bruna et al. A importância do pré-natal na prevenção de complicações materno-fetais do diabetes mellitus gestacional. *Revista Eletrônica Acervo Científico*, v. 27, p. e7588-e7588, 2021.
- ANJOS, Joyce Carolina Silva dos et al. Perfil epidemiológico das gestantes atendidas em um centro de referência em pré natal de alto risco. *Rev. para. med*, 2014.
- DE MOURA, Samilla Gonçalves et al. Assistência pré-natal realizada pelo enfermeiro (a): um olhar da mulher gestante. *Revista de pesquisa cuidado é fundamental online*, v. 7, n. 3, p. 2930-2938, 2015.